

**Supplementary Table 5. The docking results of top one results of each targets and standard drug *Baicalin*, lopinavir and arbidol.**

NO	compounds	reference	Docking (binding energy) (kJ/mol)		
			Mpro	ACE2	Spro
M1	<i>Licorice glycoside E</i>	[1]	-39.47	-31.15	-36.10
M2	<i>Xambioona</i>	[2]	-36.58	-50.50	-34.17
M3	<i>Baicalin</i>	[3, 4]	-34.66	-47.25	-31.29
M4	lopinavir	[5]	-38.99	-45.25	-29.84
M5	abidol	[5]	-26.47	-31.29	-22.14

## Supplementary References

- Nazari S, Rameshrad M, Hosseinzadeh H. Toxicological effects of *glycyrrhiza glabra* (licorice): a review. *Phytother Res.* 2017; 31:1635–50.  
<https://doi.org/10.1002/ptr.5893> PMID:[28833680](#)
- Wu C, Liu Y, Yang Y, Zhang P, Zhong W, Wang Y, Wang Q, Xu Y, Li M, Li X, Zheng M, Chen L, Li H. Analysis of therapeutic targets for SARS-CoV-2 and discovery of potential drugs by computational methods. *Acta Pharm Sin B*. 2020; 10:766–88.  
<https://doi.org/10.1016/j.apsb.2020.02.008>  
PMID:[32292689](#)
- Chen F, Chan KH, Jiang Y, Kao RY, Lu HT, Fan KW, Cheng VC, Tsui WH, Hung IF, Lee TS, Guan Y, Peiris JS, Yuen KY. *In vitro* susceptibility of 10 clinical isolates of SARS coronavirus to selected antiviral compounds. *J Clin Virol*. 2004; 31:69–75.  
<https://doi.org/10.1016/j.jcv.2004.03.003>  
PMID:[15288617](#)
- Deng YF, Aluko RE, Jin Q, Zhang Y, Yuan LJ. Inhibitory activities of baicalin against renin and angiotensin-converting enzyme. *Pharm Biol*. 2012; 50:401–06.  
<https://doi.org/10.3109/13880209.2011.608076>  
PMID:[22136493](#)
- Chen J, Ling Y, Xi X, Liu P, Li F, Li T, Shang Z, Wang M, Shen Y, Lu H. Efficacies of lopinavir/ritonavir and abidol in the treatment of novel coronavirus pneumonia. *Chinese Journal of Infectious Diseases*. 2020; 38:E008–E008.