Supplementary Material 3. Smith et al. depression criteria.

Depression (adapted from Smith et al. (2013), doi: 10.1371/journal.pone.0075362).

3. Single probable episode of major depression:

4598 ever depressed/down for a whole week; plus 4609 at least two weeks duration; plus

 $\frac{4620}{OR}$ only one episode, plus $\frac{2090}{2090}$ ever seen a GP or $\frac{2100}{2100}$ a psychiatrist for nerves, anxiety, depression OR

<u>4631</u> ever anhedonic (unenthusiasm/uninterest) for a whole week; plus <u>5375</u> at least two weeks; plus <u>5386</u> only one episode; plus 2090 ever seen a GP or 2100 a psychiatrist for nerves, anxiety, depression.

4. Probable recurrent major depression (moderate):

 $\frac{4598}{2090}$ ever depressed/down for a whole week; plus $\frac{4609}{20}$ at least two weeks duration; plus $\frac{4620}{20}$ at least two episodes; plus $\frac{2090}{2090}$ ever seen a GP (but not a psychiatrist) for nerves, anxiety, depression

OR

4631 ever anhedonic (unenthusiasm/uninterest) for a whole week; plus 5375 at least two weeks; plus

5386 at least two episodes; plus 2090 ever seen a GP (but not a psychiatrist) for nerves, anxiety, depression.

5. Probable recurrent major depression (severe):

4598 ever depressed/down for a whole week; plus 4609 at least two weeks duration; plus 4620 at least two episodes; plus 2100 ever seen a psychiatrist for nerves, anxiety, depression

OR

4631 ever anhedonic (unenthusiasm/uninterest) for a whole week; plus 5375 at least two weeks; plus

5386 at least two episodes; plus 2100 ever seen a psychiatrist for nerves, anxiety, depression.