

SUPPLEMENTARY TABLES

Supplementary Table 1. Age-specific incidence and mortality rates of Hodgkin lymphoma by age and sex in 2019 (per 100,000 population).

| Age groups (year) | Incidence | | | Mortality | | |
|-------------------|---------------------|-----------------------|---------------------|---------------------|---------------------|---------------------|
| | Both | Male | Female | Both | Male | Female |
| 0– | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| 1– | 0.18 (0.14–0.27) | 0.17 (0.11–0.29) | 0.20 (0.14–0.32) | 0.02 (0.01–0.03) | 0.02 (0.01–0.03) | 0.02 (0.02–0.03) |
| 5– | 0.24 (0.17–0.32) | 0.33 (0.23–0.48) | 0.13 (0.10–0.16) | 0.02 (0.02–0.03) | 0.03 (0.02–0.05) | 0.01 (0.01–0.02) |
| 10– | 0.17 (0.13–0.23) | 0.21 (0.15–0.32) | 0.12 (0.09–0.18) | 0.02 (0.02–0.03) | 0.02 (0.02–0.04) | 0.01 (0.01–0.02) |
| 15– | 0.46 (0.35–0.59) | 0.54 (0.36–0.73) | 0.37 (0.27–0.54) | 0.04 (0.03–0.05) | 0.05 (0.03–0.06) | 0.03 (0.02–0.05) |
| 20– | 0.63 (0.48–0.84) | 0.80 (0.52–1.07) | 0.46 (0.31–0.69) | 0.06 (0.05–0.08) | 0.07 (0.05–0.10) | 0.04 (0.03–0.07) |
| 25– | 0.48 (0.38–0.63) | 0.59 (0.40–0.80) | 0.37 (0.24–0.57) | 0.06 (0.05–0.08) | 0.07 (0.05–0.10) | 0.05 (0.03–0.07) |
| 30– | 0.56 (0.42–0.71) | 0.77 (0.49–1.01) | 0.35 (0.24–0.53) | 0.08 (0.06–0.10) | 0.10 (0.07–0.14) | 0.05 (0.04–0.07) |
| 35– | 0.47 (0.35–0.60) | 0.63 (0.41–0.85) | 0.31 (0.21–0.46) | 0.08 (0.06–0.10) | 0.10 (0.07–0.14) | 0.05 (0.04–0.07) |
| 40– | 0.53 (0.38–0.67) | 0.70 (0.46–0.95) | 0.35 (0.23–0.49) | 0.10 (0.08–0.13) | 0.13 (0.09–0.18) | 0.07 (0.05–0.09) |
| 45– | 0.53 (0.38–0.68) | 0.73 (0.48–1.01) | 0.31 (0.20–0.43) | 0.11 (0.09–0.15) | 0.16 (0.11–0.23) | 0.07 (0.05–0.09) |
| 50– | 0.73 (0.52–0.92) | 0.95 (0.60–1.30) | 0.50 (0.33–0.69) | 0.27 (0.20–0.34) | 0.23 (0.15–0.31) | 0.12 (0.08–0.16) |
| 55– | 0.83 (0.59–1.06) | 1.14 (0.74–1.57) | 0.52 (0.33–0.72) | 0.37 (0.27–0.47) | 0.36 (0.24–0.50) | 0.17 (0.11–0.23) |
| 60– | 1.05 (0.73–1.33) | 1.46 (0.90–1.95) | 0.64 (0.38–0.87) | 0.54 (0.39–0.67) | 0.52 (0.36–0.70) | 0.23 (0.15–0.31) |
| 65– | 1.19 (0.83–1.49) | 1.61 (0.97–2.14) | 0.79 (0.50–1.08) | 0.74 (0.54–0.90) | 0.73 (0.47–0.99) | 0.37 (0.24–0.49) |
| 70– | 1.33 (0.91–1.63) | 1.80 (1.09–2.37) | 0.87 (0.53–1.16) | 0.91 (0.70–1.23) | 1.00 (0.64–1.32) | 0.50 (0.32–0.66) |
| 75– | 1.63 (1.07–2.01) | 2.10 (1.21–2.72) | 1.21 (0.73–1.62) | 0.91 (0.63–1.12) | 1.16 (0.71–1.53) | 0.69 (0.43–0.90) |
| 80– | 2.19 (1.38–2.72) | 2.95 (1.63–3.90) | 1.59 (0.94–2.14) | 1.05 (0.69–1.28) | 1.39 (0.81–1.80) | 0.78 (0.48–1.04) |
| 85– | 2.63 (1.63–3.29) | 4.47 (2.39–5.82) | 1.64 (0.92–2.20) | 1.26 (0.82–1.52) | 2.10 (1.21–2.69) | 0.80 (0.48–1.07) |
| 90– | 5.99 (3.25–8.28) | 13.21 (6.15–19.23) | 4.49 (2.22–6.66) | 0.86 (0.50–1.13) | 1.76 (0.86–2.40) | 0.68 (0.35–0.94) |
| 95– | 4.92 (2.38–7.44) | 11.98 (5.25–20.89) | 4.30 (1.91–6.68) | 0.81 (0.41–1.18) | 1.31 (0.60–1.93) | 0.77 (0.37–1.14) |

Data in parentheses are 95% uncertainty intervals.

Supplementary Table 2. Age-specific incidence and mortality rates of non-Hodgkin lymphoma by age and sex in 2019 (per 100,000 population).

| Age groups (year) | Incidence | | | Mortality | | |
|-------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|
| | Both | Male | Female | Both | Male | Female |
| 0– | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| 1– | 0.00 | 0.00 | 0.00 | 0.28 (0.22–0.37) | 0.31 (0.23–0.41) | 0.26 (0.21–0.33) |
| 5– | 1.77 (1.44–2.21) | 2.28 (1.78–2.95) | 1.18 (0.98–1.45) | 0.29 (0.24–0.36) | 0.38 (0.30–0.48) | 0.19 (0.16–0.23) |
| 10– | 1.53 (1.28–1.84) | 1.81 (1.43–2.31) | 1.21 (1.00–1.44) | 0.26 (0.22–0.30) | 0.31 (0.25–0.38) | 0.20 (0.17–0.24) |
| 15– | 2.24 (1.85–2.70) | 2.79 (2.14–3.60) | 1.62 (1.36–1.95) | 0.39 (0.32–0.46) | 0.49 (0.37–0.61) | 0.27 (0.23–0.32) |
| 20– | 1.81 (1.48–2.24) | 2.31 (1.79–2.97) | 1.27 (0.97–1.65) | 0.53 (0.44–0.64) | 0.68 (0.54–0.87) | 0.35 (0.27–0.45) |
| 25– | 1.88 (1.56–2.29) | 2.45 (1.97–3.06) | 1.29 (0.94–1.71) | 0.55 (0.47–0.65) | 0.72 (0.59–0.88) | 0.37 (0.28–0.48) |
| 30– | 2.40 (2.00–2.93) | 3.16 (2.54–3.94) | 1.63 (1.22–2.15) | 0.72 (0.61–0.84) | 0.96 (0.78–1.17) | 0.48 (0.36–0.61) |
| 35– | 3.24 (2.70–3.95) | 4.22 (3.32–5.36) | 2.23 (1.67–2.89) | 1.00 (0.84–1.18) | 1.30 (1.04–1.61) | 0.68 (0.51–0.86) |
| 40– | 3.01 (2.44–3.68) | 3.99 (3.08–5.15) | 1.99 (1.53–2.55) | 1.43 (1.18–1.71) | 1.92 (1.49–2.41) | 0.93 (0.71–1.17) |
| 45– | 4.15 (3.30–5.16) | 5.81 (4.35–7.70) | 2.43 (1.85–3.12) | 2.02 (1.64–2.47) | 2.85 (2.15–3.69) | 1.16 (0.90–1.46) |
| 50– | 6.59 (5.23–8.18) | 9.24 (6.87–12.20) | 3.91 (2.99–5.03) | 3.22 (2.60–3.94) | 4.55 (3.42–5.85) | 1.88 (1.46–2.40) |
| 55– | 9.98 (7.93–12.35) | 13.91 (10.39–18.30) | 6.00 (4.63–7.67) | 4.87 (3.93–5.96) | 6.83 (5.17–8.75) | 2.89 (2.26–3.67) |
| 60– | 12.76 (10.31–15.49) | 17.06 (12.92–21.82) | 8.43 (6.55–10.72) | 6.51 (5.41–7.77) | 8.73 (6.75–11.08) | 4.27 (3.38–5.33) |
| 65– | 18.14 (14.89–22.02) | 24.14 (18.64–30.52) | 12.34 (9.81–15.65) | 9.25 (7.75–10.86) | 12.34 (9.76–15.39) | 6.26 (5.00–7.69) |
| 70– | 25.70 (21.06–30.95) | 34.43 (26.73–43.40) | 17.39 (13.81–21.99) | 13.08 (11.11–15.43) | 17.55 (13.96–21.90) | 8.82 (7.14–10.76) |
| 75– | 29.09 (24.21–34.64) | 37.95 (30.31–46.80) | 21.11 (16.65–26.37) | 14.78 (12.61–17.21) | 19.26 (15.61–23.61) | 10.74 (8.71–13.06) |
| 80– | 25.42 (21.45–30.02) | 34.37 (27.92–41.69) | 18.42 (14.42–22.77) | 17.13 (14.66–19.76) | 23.09 (19.03–27.68) | 12.46 (10.04–14.92) |
| 85– | 32.75 (27.51–38.41) | 54.62 (45.78–63.72) | 20.97 (16.21–26.45) | 21.82 (18.70–24.78) | 36.47 (31.13–42.12) | 13.93 (10.86–16.98) |
| 90– | 28.04 (22.16–33.24) | 54.21 (44.12–65.19) | 22.60 (17.13–27.90) | 18.39 (14.96–21.36) | 35.61 (29.59–41.70) | 14.81 (11.27–17.86) |
| 95– | 28.52 (21.26–35.73) | 49.97 (40.55–59.70) | 26.63 (19.08–34.22) | 18.66 (14.07–22.44) | 31.02 (25.12–36.64) | 17.57 (12.71–21.54) |

Data in parentheses are 95% uncertainty intervals.

Supplementary Table 3. Age-standardized incidence, mortality and prevalence rates of lymphoma by province of China in 2019 (per 100,000 population).

| Province | Hodgkin lymphoma | | | Non-Hodgkin lymphoma | | |
|----------------|---------------------|---------------------|---------------------|----------------------|---------------------|------------------------|
| | Incidence | Mortality | Prevalence | Incidence | Mortality | Prevalence |
| Anhui | 0.58 (0.41–0.75) | 0.15 (0.10–0.19) | 3.95 (2.83–5.13) | 5.67 (4.44–7.07) | 2.61 (2.07–3.23) | 26.55 (19.82–34.92) |
| Beijing | 0.76 (0.52–0.97) | 0.11 (0.07–0.14) | 5.96 (4.14–7.68) | 6.29 (5.04–7.62) | 2.38 (1.93–2.88) | 35.36 (26.87–44.82) |
| Chongqing | 0.51 (0.37–0.80) | 0.12 (0.09–0.19) | 3.60 (2.54–5.62) | 5.34 (4.15–6.81) | 2.41 (1.89–3.04) | 25.68 (18.89–34.30) |
| Fujian | 0.60 (0.45–0.78) | 0.13 (0.09–0.16) | 4.30 (3.22–5.76) | 5.35 (4.33–6.63) | 2.33 (1.89–2.84) | 26.37 (19.83–34.13) |
| Gansu | 0.30 (0.21–0.50) | 0.13 (0.09–0.22) | 1.57 (1.04–2.70) | 3.52 (2.87–4.36) | 2.02 (1.65–2.43) | 9.48 (6.50–13.03) |
| Guangdong | 0.52 (0.39–0.68) | 0.11 (0.08–0.14) | 3.81 (2.85–4.99) | 3.87 (3.12–4.77) | 1.66 (1.37–2.00) | 16.65 (12.06–22.04) |
| Guangxi | 0.44 (0.33–0.61) | 0.16 (0.12–0.21) | 2.59 (1.91–3.61) | 4.39 (3.44–5.44) | 2.29 (1.78–2.87) | 15.64 (10.88–21.22) |
| Guizhou | 0.24 (0.14–0.49) | 0.13 (0.07–0.25) | 1.06 (0.58–2.22) | 3.67 (2.76–4.81) | 2.35 (1.80–3.06) | 7.62 (4.52–11.14) |
| Hainan | 0.33 (0.20–0.56) | 0.12 (0.08–0.19) | 1.94 (1.19–3.42) | 4.11 (3.16–5.17) | 2.16 (1.71–2.69) | 14.50 (10.14–19.56) |
| Hebei | 0.73 (0.41–0.99) | 0.21 (0.12–0.29) | 4.79 (2.69–6.63) | 5.64 (4.54–6.98) | 2.67 (2.14–3.23) | 23.01 (16.85–30.72) |
| Heilongjiang | 0.59 (0.43–0.78) | 0.16 (0.11–0.21) | 3.92 (2.87–5.31) | 4.59 (3.69–5.69) | 2.16 (1.77–2.63) | 19.35 (14.20–25.59) |
| Henan | 0.52 (0.39–0.72) | 0.13 (0.10–0.18) | 3.50 (2.61–5.00) | 4.04 (3.28–5.05) | 1.85 (1.48–2.28) | 16.53 (11.57–22.26) |
| Hubei | 0.57 (0.40–0.75) | 0.15 (0.10–0.20) | 3.86 (2.73–5.12) | 5.26 (4.15–6.71) | 2.45 (1.96–3.05) | 23.79 (17.66–31.24) |
| Hunan | 0.41 (0.30–0.62) | 0.13 (0.10–0.20) | 2.49 (1.81–3.85) | 5.65 (4.61–7.00) | 2.84 (2.27–3.49) | 23.67 (17.79–30.94) |
| Inner Mongolia | 0.44 (0.31–0.73) | 0.12 (0.08–0.19) | 2.98 (2.04–5.03) | 5.51 (4.40–6.82) | 2.59 (2.07–3.19) | 24.63 (18.56–32.53) |
| Jiangsu | 0.80 (0.48–1.10) | 0.15 (0.08–0.21) | 5.97 (3.62–8.16) | 5.92 (4.59–7.55) | 2.45 (1.92–3.07) | 30.52 (22.58–39.83) |
| Jiangxi | 0.46 (0.34–0.58) | 0.16 (0.12–0.20) | 2.71 (2.02–3.58) | 4.17 (3.36–5.12) | 2.14 (1.75–2.60) | 15.77 (11.43–20.97) |
| Jilin | 0.55 (0.41–0.71) | 0.14 (0.10–0.17) | 3.81 (2.80–5.04) | 5.14 (4.26–6.35) | 2.31 (1.98–2.81) | 24.38 (18.72–32.03) |
| Liaoning | 0.83 (0.49–1.10) | 0.18 (0.10–0.24) | 6.02 (3.60–7.91) | 6.67 (5.37–8.27) | 2.86 (2.32–3.53) | 34.85 (26.59–45.51) |
| Ningxia | 0.41 (0.29–0.66) | 0.13 (0.09–0.20) | 2.63 (1.80–4.37) | 4.25 (3.34–5.27) | 2.10 (1.64–2.62) | 15.87 (11.49–21.08) |
| Qinghai | 0.25 (0.14–0.51) | 0.14 (0.08–0.29) | 0.98 (0.51–2.09) | 3.74 (2.94–4.68) | 2.51 (1.98–3.08) | 6.63 (4.08–9.84) |
| Shaanxi | 0.40 (0.28–0.62) | 0.13 (0.09–0.19) | 2.53 (1.71–4.03) | 4.33 (3.35–5.52) | 2.16 (1.66–2.68) | 17.11 (11.99–23.12) |
| Shandong | 0.67 (0.48–0.85) | 0.13 (0.09–0.16) | 4.92 (3.54–6.32) | 4.48 (3.57–5.55) | 1.85 (1.50–2.26) | 21.42 (15.37–27.65) |
| Shanghai | 0.63 (0.45–0.84) | 0.10 (0.07–0.13) | 4.88 (3.54–6.55) | 6.87 (5.49–8.40) | 2.75 (2.21–3.33) | 38.57 (29.69–49.20) |
| Shanxi | 0.46 (0.32–0.72) | 0.12 (0.09–0.19) | 3.12 (2.11–4.99) | 5.44 (4.19–6.74) | 2.55 (1.96–3.13) | 24.69 (18.03–32.43) |

| | | | | | | |
|------------|---------------------|---------------------|-----------------------|---------------------|---------------------|------------------------|
| Sichuan | 0.49 (0.36–0.67) | 0.15 (0.12–0.21) | 3.07 (2.24–4.36) | 5.55 (4.38–6.95) | 2.76 (2.22–3.43) | 22.90 (16.77–30.97) |
| Tianjin | 0.68 (0.49–0.88) | 0.12 (0.08–0.15) | 5.15 (3.81–6.75) | 6.73 (5.39–8.30) | 2.73 (2.18–3.29) | 36.88 (27.86–46.98) |
| Tibet | 0.25 (0.14–0.51) | 0.20 (0.11–0.40) | 0.53 (0.29–1.08) | 3.01 (2.34–3.81) | 2.72 (2.11–3.39) | 2.12 (0.73–3.93) |
| Xinjiang | 0.50 (0.32–0.66) | 0.26 (0.17–0.34) | 2.19 (1.37–3.00) | 3.90 (3.20–4.86) | 2.45 (2.00–3.02) | 8.78 (5.72–12.55) |
| Yunnan | 0.30 (0.20–0.52) | 0.15 (0.10–0.25) | 1.39 (0.89–2.46) | 3.52 (2.85–4.33) | 2.17 (1.78–2.65) | 8.58 (5.68–11.99) |
| Zhejiang | 1.02 (0.50–1.45) | 0.18 (0.08–0.25) | 7.71 (3.87–10.92) | 4.68 (3.82–5.73) | 1.89 (1.52–2.29) | 23.17 (17.39–30.12) |
| Hong Kong* | 1.10 (0.65–1.54) | 0.11 (0.07–0.15) | 8.83 (5.23–12.51) | 5.25 (3.94–7.03) | 2.13 (1.60–2.83) | 27.69 (19.36–37.78) |
| Macao* | 1.29 (0.59–1.94) | 0.13 (0.06–0.20) | 10.36 (4.76–15.60) | 4.17 (3.09–5.60) | 1.63 (1.21–2.18) | 21.13 (14.69–29.02) |

*Special administrative regions. Data in parentheses are 95% uncertainty intervals.